

- Electronic copy is controlled under document control procedure. Hard copy is uncontrolled & under responsibility of beholder. • النسخة الإلكترونية هي النسخة المضبوطة وفق إجراء ضبط الوثائق- النسخ الورقية غير مضبوطة وتقع على مسؤولية حاملها.
- It is allowed ONLY to access and keep this document with who issued, who is responsible and to whom it is applicable. • يسمح بالوصول وباحتفاظ بهذه الوثيقة مع مصدرها أو مع المسؤول عن تطبيقها أو مع المطبق عليهم.
- Information security code:  Open  Shared -Confidential  Shared-Sensitive  Shared-Secret  مشارك -مشارك  بيانات مفتوحة  مشارك -خاص  مشارك -حساس  مشارك -سري

# DHA TELEHEALTH CLINICAL GUIDELINES

## FOR VIRTUAL MANAGEMENT

### OF MINOR HEAD INJURY – 31

#### Version 2

Issue date: 21/02/2024

Effective date: 21/04/2024

Health Policies and Standards Department

Health Regulation Sector (2024)

## INTRODUCTION

Health Regulation Sector (HRS) forms an integral part of Dubai Health Authority (DHA) and is mandated by DHA Law No. (14) of the year (2021) amending some clauses of law No. (6) of 2018 pertaining to the Dubai Health Authority (DHA), to undertake several functions including but not limited to:

- Developing regulation, policy, standards, guidelines to improve quality and patient safety and promote the growth and development of the health sector;
- Licensure and inspection of health facilities as well as healthcare professionals and ensuring compliance to best practice;
- Managing patient complaints and assuring patient and physician rights are upheld;
- Governing the use of narcotics, controlled and semi-controlled medications;
- Strengthening health tourism and assuring ongoing growth; and
- Assuring management of health informatics, e-health and promoting innovation.

The DHA Telehealth Clinical Guidelines aim to fulfil the following overarching DHA Strategic Priorities (2026):

- Pioneering Human-centered health system to promote trust, safety, quality and care for patients and their families.
- Make Dubai a lighthouse for healthcare governance, integration and regulation.
- Leading global efforts to combat epidemics and infectious diseases and prepare for disasters.

- Pioneering prevention efforts against non-communicable diseases.
- Become a global digital health hub.
- Foster healthcare education, research and innovation.

## ACKNOWLEDGMENT

The Health Policy and Standards Department (HPSD) developed this Guideline in collaboration with Subject Matter Experts and would like to acknowledge and thank these health professionals for their dedication toward improving quality and safety of healthcare services in the Emirate of Dubai.

### Health Regulation Sector

#### Dubai Health Authority

## TABLE OF CONTENTS

|   |           |
|---|-----------|
| <b>EXECUTIVE SUMMARY</b>  | <b>5</b>  |
| <b>DEFINITIONS/ABBREVIATIONS</b>                                      | <b>6</b>  |
| <b>1. BACKGROUND</b>  | <b>7</b>  |
| <b>2. SCOPE</b>   | <b>7</b>  |
| <b>3. PURPOSE</b>   | <b>7</b>  |
| <b>4. APPLICABILITY</b>   | <b>8</b>  |
| <b>5. RECOMMENDATION</b>  | <b>8</b>  |
| <b>6. RED FLAGS</b>   | <b>9</b>  |
| <b>7. DIFFERENTIAL DIAGNOSIS</b>                                      | <b>11</b> |
| <b>8. INVESTIGATIONS</b>  | <b>12</b> |
| <b>9. REFERRAL CRITERIA</b>   | <b>12</b> |
| <b>10. MANAGEMENT</b>   | <b>13</b> |
| <b>REFERENCES</b>   | <b>18</b> |
| <b>APPENDIX 1 – VIRTUAL MANAGEMENT OF MINOR HEAD INJURY ALGORITHM</b> | <b>19</b> |

## EXECUTIVE SUMMARY

Telehealth is based on Evidence Based Practice (EBP) which is the conscientious, explicit and judicious use of current best evidence in making decisions about the care of the individual patient.

It means integrating individual clinical expertise with the best available external clinical evidence and guidelines from systematic research.

EBP is important because it aims to provide the most effective care virtually, with the aim of improving patient outcomes. As health professionals, part of providing a professional service is ensuring that practice is informed by the best available evidence.

This guideline is presented in the format comprising of clinical history/symptoms, differential diagnosis, investigations and management. Identification of 'Red Flags' or serious conditions associated with the disease is an essential part of this telehealth guideline as it aids the physician to manage patients safely and appropriately by referrals to ER, family physicians or specialists for a face to face management.

Minor head injuries are common in people of all ages and rarely result in any permanent brain damage. The most common causes of head injuries are falls, assaults, and road traffic collisions. Each year millions of adults and children attend emergency departments with a head injury all over the world but more than 80% have only a minor injury.

Children are more likely to sustain a minor head injury because they're very active.

The symptoms of mild head injury are usually mild and short lived. They may include a minor headache, nausea, mild dizziness, mild blurred vision.

## DEFINITIONS/ABBREVIATIONS

**Virtual Clinical Assessment:** Is the evaluation of the patient's medical condition virtually via telephone or video call consultations, which may include one or more of the following: patient medical history, physical examination and diagnostic investigations.

**Patient:** The person who receives the healthcare services or the medical investigation or treatment provided by a DHA licensed healthcare professional.

## ABBREVIATIONS

|            |   |                                      |
|------------|---|--------------------------------------|
| <b>DHA</b> | : | Dubai Health Authority               |
| <b>EBP</b> | : | Evidence Based Practice              |
| <b>ER</b>  | : | Emergency Room                       |
| <b>HRS</b> | : | Health Regulation Sector             |
| <b>EBP</b> | : | Nonsteroidal Anti-Inflammatory Drugs |
| <b>TBI</b> | : | Traumatic Brain Injuries             |

## 1. BACKGROUND

1.1. In general, head injuries can be divided into two categories based on what causes them. They can either be head injuries due to blows to the head or head injuries due to shaking.

1.1.1. Head injuries caused by a blow to the head are usually associated with:

- a. Motor vehicle accidents
- b. Falls
- c. Physical assaults
- d. Sports related accidents

1.2. In most cases, skull usually protect the brain from serious harm. However, injuries severe enough to cause head injury can also be associated with injuries to the brain and spinal cord.

1.3. Head injuries caused by shaking are most common in infants and young children but they can occur any time you experience violent shaking.

## 2. SCOPE

2.1. Telehealth services in DHA licensed Health Facilities.

## 3. PURPOSE

3.1. To support the implementation of Telehealth services for patients with complaints of Minor Head Injury in Dubai Health Authority (DHA) licensed Health Facilities

#### 4. APPLICABILITY

- 4.1. DHA licensed physicians and health facilities providing Telehealth services.
- 4.2. Exclusion for Telehealth services are as follows
  - 4.2.1. Emergency cases where immediate intervention or referral is required
  - 4.2.2. Prescribe Narcotics, Controlled or Semi-Controlled medications

#### 5. RECOMMENDATION

##### 5.1. Virtual Clinical Assessment

##### 5.1.1. Signs of minor head injuries

Often cause a bump or bruise. As long as the person is conscious and with no deep wounds, it's unlikely there will be any serious damage. Other symptoms of a minor head injury may include:

- a. A mild headache
- b. Nausea (feeling sick)
- c. Mild dizziness
- d. Mild blurred vision

##### 5.1.2. Signs of serious head injury

- a. Unconsciousness, either briefly or for a longer period of time
- b. Difficulty staying awake or still being sleepy several hours after the injury

- c. Clear fluid leaking from the nose or ears – this could be cerebrospinal fluid, which normally surrounds the brain
- d. Bleeding from one or both ears
- e. Bruising behind one or both ears
- f. Any sign of skull damage or a penetrating head injury
- g. Difficulty speaking, such as slurred speech
- h. Difficulty understanding what people say
- i. Reading or writing problems
- j. Balance problems or difficulty walking
- k. Loss of power or sensation in part of the body, such as arm or leg
- l. General weakness
- m. Vision problems, such as significantly blurred or double vision
- n. Having a seizure or fit
- o. Memory loss amnesia
- p. Persistent headache
- q. Vomiting since the injury
- r. Irritability or unusual behavior

## 6. RED FLAGS

While assessing a patient with minor head injury the following red flags symptoms must be ruled out:

- 6.1. Unconsciousness – either brief concussion or for a longer period of time
- 6.2. Fits or seizures
- 6.3. Problems with the senses – such as hearing loss or double vision
- 6.4. Repeated vomiting
- 6.5. Blood or clear fluid coming from the ears or nose
- 6.6. Bruising behind one or both ears
- 6.7. Memory loss amnesia
- 6.8. The injury was caused by a forceful blow to the head at speed
- 6.9. The person had brain surgery before
- 6.10. The person has had problems with uncontrollable bleeding or a blood clotting disorder, or is taking medication that may cause bleeding problems, such as Warfarin
- 6.11. The person is intoxicated by drugs or alcohol
- 6.12. It is possible the injury wasn't accidental – for example, child abuse and Munchausen syndrome by proxy
- 6.13. Irritability or unusual behavior
- 6.14. Difficulty staying awake or still being sleepy several hours after the injury
- 6.15. Any sign of skull damage or a penetrating head injury
- 6.16. Difficulty speaking or understanding what people say
- 6.17. Balance problems or difficulty walking
- 6.18. Loss of power or sensation in part of the body, such as arm or leg

6.19. Significantly blurred or double vision

6.20. Persistent headache

## 7. DIFFERENTIAL DIAGNOSIS

7.1. Most of the patients with mild traumatic brain injuries (TBI) recover full function within about 3 months, but a significant minority do not. Failure to recover as expected following a diagnosed or suspected mild TBI is most commonly related to a concurrent diagnosis or alternative diagnosis or condition. Consideration during the diagnostic process must be given to

7.1.1. Alternative organic conditions (prior or unsuspected severe TBI, pain, medication side effects or dementia)

7.1.2. Pre-existing non-organic conditions (active or dormant psychiatric conditions)

7.1.3. Pre-existing personality characteristics, or social/economic factors

7.1.4. Non-clinical conditions (compensation/litigation, malingering)

7.1.5. Post-injury psychiatric morbidity (notably depression, anxiety, post-traumatic stress disorder, panic disorder, or conversion disorder)

7.2. Rigorous care must be given during the clinical assessment of the patient to evaluate for different or contributory diagnostic possibilities.

- 7.3. Not all patients that have symptoms following a blow to the head have traumatic brain injury as the sole etiologic agent for their symptoms. Accurate diagnosis will lead to better treatment and optimal outcomes.

## 8. INVESTIGATIONS

- 8.1. Based on assessment, this will include CT Scan, MRI, X-ray skull.

## 9. REFERRAL CRITERIA

- 9.1. Seek immediate medical attention if, after a knock to the head, you notice any of these symptoms:

- 9.1.1. Unconsciousness – either brief concussion or for a longer period of time
- 9.1.2. Fits or seizures
- 9.1.3. Problems with the senses – such as hearing loss or double vision
- 9.1.4. Repeated vomiting
- 9.1.5. Blood or clear fluid coming from the ears or nose
- 9.1.6. Bruising behind one or both ears
- 9.1.7. Memory loss amnesia
- 9.1.8. The injury was caused by a forceful blow to the head at speed
- 9.1.9. The person had brain surgery before
- 9.1.10. The person has had problems with uncontrollable bleeding or a blood clotting disorder, or is taking medication that may cause bleeding problems, such as Warfarin

- 9.1.11. The person is intoxicated by drugs or alcohol
- 9.1.12. It is possible the injury wasn't accidental – for example, child abuse and Munchausen syndrome by proxy
- 9.1.13. Irritability or unusual behaviour
- 9.1.14. Difficulty staying awake or still being sleepy several hours after the injury
- 9.1.15. Any sign of skull damage or a penetrating head injury
- 9.1.16. Difficulty speaking or understanding what people say
- 9.1.17. Balance problems or difficulty walking
- 9.1.18. Loss of power or sensation in part of the body, such as arm or leg
- 9.1.19. Significantly blurred or double vision
- 9.1.20. Persistent headache
- 9.1.21. If symptoms persist two weeks after the head injury
- 9.2. If any of these symptoms are present, particularly a loss of consciousness – even if only for a short period of time – refer immediately to emergency department or call for an ambulance.

## 10. MANAGEMENT

- 10.1. Refer to APPENDIX 1 for the Virtual Management of Minor Head Injury Algorithm
- 10.2. Treating a minor head injury
  - 10.2.1. If the patient experiences a knock, bump or blow to the head, advise the patient/caregiver to:

- a. Sit down, comfort the patient, and make sure they rest.
- b. Hold a cold compress to their head – try a bag of ice or frozen peas wrapped in a tea towel.
- c. For the first 24 hours after the injury, it's important for someone to stay with the injured person to keep an eye out for any new symptoms that develop.
- d. It is also important to advise the patient to rest, avoid aggravating the injury with stressful situations, and avoid contact sports until fully recovered.
- e. Mild headaches can be treated with paracetamol
- f. Seek immediate medical advice if symptoms such as dizziness and a headache get worse.

10.2.2. Most people who attend hospital with a minor head injury are allowed to return home shortly afterwards and will make a full recovery within a few days.

### 10.3. Advice for adults

10.3.1. If a patient sustained a minor head injury:

- a. Patient should not drink alcohol or take recreational drugs
- b. Patient should not take sleeping pills, sedatives or tranquillizers
- c. Take paracetamol or NSAID if have a mild headache

- d. Patient should not play contact sport, such as football, for at least 3 weeks
- e. Patient should not return to work, college or school until completely recovered and feel ready
- f. Patient should not drive a car, motorbike or bicycle or operate machinery until completely recovered

#### 10.4. Advice for children

##### 10.4.1. If a child has got minor head injury:

- a. Give them paracetamol or NSAIDs if they have a mild headache
- b. Avoid getting them too excited
- c. Don't have too many visitors
- d. Don't let them play contact sports, such as football for at least 3 weeks
- e. Make sure they avoid rough play for a few days

#### 10.5. Prevention

10.5.1. Many head injuries are the result of accidents that are difficult to predict or avoid. But there are ways to reduce the risk.

#### 10.6. Safety helmets

10.6.1. Cyclists and motorcyclists can protect their heads by wearing a properly fitting safety helmet

- a. As well as wearing a helmet when cycling, one should also make sure:
- b. Use lights and wear reflective clothing when cycling in the dark
- c. Are aware of the dangers of the road and know how to stay safe
- d. Check bikes are in good working order

## 10.7. Safety at home

10.7.1. Following sensible health and safety guidelines can help prevent accidents at home. Advice that will help keep home and garden as safe as possible includes:

- a. Keeping stairways tidy so you don't trip over anything
- b. Not standing on an unstable chair to change a light bulb – use a stepladder instead
- c. Cleaning up any spillages to prevent someone slipping over

## 10.8. Childproofing the home

10.8.1. It's not possible to childproof all home completely. But one can take steps to keep toddlers and young children safe at home:

- a. Check windows are lockable and can't be opened by your child, particularly bedroom windows
- b. Move furniture, such as beds, sofas and chairs, away from windows to prevent child climbing up and falling out
- c. Fit safety gates at the top and bottom of stairs

## 10.9. Safety at work

10.9.1. To reduce the risk of sustaining a head injury at work, always follow the following:

- a. Wear a hard hat when working in potentially hazardous areas.
- b. Only use ladders in a workplace environment for short-term light work.
- c. Any work that requires spending a considerable amount of time at height or involves heavy lifting should be carried out on scaffolding or another suitable platform
- d. Any work that involves going up on to a roof should also be considered high risk, and high standards of safety are therefore essential.

## 10.10. Sport safety

10.10.1. Advise the patient to wear any necessary safety equipment when playing sports, particularly contact sports.

## REFERENCES

1. NHS forum. [Internet]. Minor Head injury. [updated 2021; cited 2022 May 24]. Available from:  
<https://www.nhsinform.scot/illnesses-and-conditions/injuries/head-and-neck-injuries/minor-head-injury>
2. Rospa.com. (2019). *The Royal Society for the Prevention of Accidents - RoSPA*. [online] Available at:  
<https://www.rospace.com/> [Accessed 11 May. 2023].
3. WILBERGER.JAMES (2019). *Overview of Head Injuries*. [online] Merck Manuals Consumer Version. Available at: <https://www.merckmanuals.com/home/injuries-and-poisoning/head-injuries/overview-of-head-injuries> [Accessed 11 May. 2023].

## APPENDIX 1 – VIRTUAL MANAGEMENT OF MINOR HEAD INJURY ALGORITHM

